

Diagnosed with NASH with liver scarring?

A guide for talking with your liver specialist

1. I have NASH with liver scarring; how can I check the state of my liver scarring?

Testing is important to help check the level of scarring on your liver. There are different types of tests your liver specialist may recommend, like:

- Blood-based tests
- Imaging and scanning tests
- Liver biopsy

If left untreated, NASH can lead to even more serious conditions like cirrhosis, cancer, or the need for a liver transplant.

Notes _____

2. Can Rezdiffra help reduce my NASH with liver scarring?

Rezdiffra is the **FIRST AND ONLY** FDA-approved treatment for NASH with liver scarring. It works directly in the liver to help reduce liver scarring and NASH. Here are some tips that can help when taking Rezdiffra:

- Take Rezdiffra once a day, every day, with or without food
- Keep taking Rezdiffra as prescribed for as long as your liver specialist tells you to

Continued on next page →

Notes _____

INDICATION AND IMPORTANT SAFETY INFORMATION

What is Rezdiffra?

Rezdiffra is a prescribed medicine used along with diet and exercise to treat adults with nonalcoholic steatohepatitis (NASH) with moderate to advanced liver scarring (fibrosis), but not with cirrhosis of the liver.

This indication is approved based on improvement of NASH and liver scarring (fibrosis). There are ongoing studies to confirm the clinical benefit of Rezdiffra.

Before you take Rezdiffra, tell your healthcare provider about all of your medical conditions, including if you:

- have any liver problems other than NASH.
- have gallbladder problems or have been told you have gallbladder problems, including gallstones.
- are pregnant or plan to become pregnant. It is not known if Rezdiffra will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if Rezdiffra passes into your breast milk. Talk to your

healthcare provider about the best way to feed your baby if you take Rezdiffra.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

- Rezdiffra and other medicines may affect each other, causing side effects. Rezdiffra may affect the way other medicines work, and other medicines may affect how Rezdiffra works.
- Especially tell your healthcare provider if you take medicines that contain gemfibrozil to help lower your triglycerides, or cyclosporine to suppress your immune system, because Rezdiffra is not recommended in patients taking these medicines.
- Tell your healthcare provider if you are taking medicines such as clopidogrel to thin your blood or statin medicines to help lower your cholesterol.

Please see the full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.

3. What are possible side effects of Rezdiffra?

Rezdiffra may cause serious side effects. Stop taking Rezdiffra and call your doctor if you develop symptoms of:

- **Liver injury (1 case in clinical trials):** including tiredness, nausea, vomiting, fever, rash, jaundice, or stomach pain/tenderness
- **Gallstones or gallbladder/pancreas inflammation:** including nausea, vomiting, fever, or stomach pain

Common side effects observed in studies include diarrhea, nausea, itching, stomach pain, vomiting, dizziness, and constipation.

See Important Safety Information below for complete list of side effects.

Diarrhea and nausea were the most common side effects. Both typically began early in treatment and were mild to moderate, lasting ~3 to 4 weeks.*

*Should these problems not resolve or should they become worse, please contact your doctor.

INDICATION AND IMPORTANT SAFETY INFORMATION (cont.)

- Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

What are the possible side effects of Rezdiffra?

Rezdiffra may cause serious side effects, including:

- **liver injury (hepatotoxicity).** Stop taking Rezdiffra and call your healthcare provider right away if you develop the following signs or symptoms of hepatotoxicity, including tiredness, nausea, vomiting, fever, rash, your skin or the white part of your eyes turns yellow (jaundice), or stomach pain/tenderness.
- **gallbladder problems.** Gallbladder problems such as gallstones, or inflammation of the gallbladder, or inflammation of the pancreas from gallstones can occur with NASH and may occur if you take Rezdiffra. Call your healthcare provider right away if you develop any signs or symptoms of these conditions, including nausea, vomiting, fever, or pain in your stomach area (abdomen)

4. How can I get Rezdiffra?

Rezdiffra is sent to you through a specialty pharmacy in Madrigal's distribution network or through the Madrigal Patient Support program. Here's how it works:

- Your liver specialist will send your prescription to a specialty pharmacy
- The specialty pharmacy will confirm insurance coverage and out-of-pocket cost
- The specialty pharmacy will call you to schedule your Rezdiffra shipment
- When it's time for a refill, the specialty pharmacy will give you a call

Notes _____

that is severe and will not go away. The pain may be felt going from your abdomen to your back and the pain may happen with or without vomiting.

The most common side effects of Rezdiffra include:

diarrhea, nausea, itching, stomach pain, vomiting, dizziness, and constipation.

These are not all the possible side effects of Rezdiffra. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at **1-800-FDA-1088** or www.fda.gov/medwatch. You may also report side effects to Madrigal Pharmaceuticals, Inc. at 1-800-905-0324.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.