

Rezdiffra™
resmetirom tablets
60mg - 80mg - 100mg

For adults with NASH with moderate to advanced liver scarring (fibrosis) without cirrhosis.

The **FIRST** FDA-approved treatment for NASH with liver scarring, along with diet and exercise, is here.

Help reduce liver scarring and NASH.

Find your way forward.

NASH=nonalcoholic steatohepatitis. NASH is also known as MAFLD, or metabolic dysfunction-associated steatohepatitis.



INDICATION AND IMPORTANT SAFETY INFORMATION

What is Rezdiffra?

Rezdiffra is a prescribed medicine used along with diet and exercise to treat adults with nonalcoholic steatohepatitis (NASH) with moderate to advanced liver scarring (fibrosis), but not with cirrhosis of the liver.


This indication is approved based on improvement of

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.

NASH and liver scarring (fibrosis). There are ongoing studies to confirm the clinical benefit of Rezdiffra.

Before you take Rezdiffra, tell your healthcare provider about all of your medical conditions, including if you:

- have any liver problems other than NASH.



Welcome!
Let's get started.

Table of Contents

ABOUT NASH

- **Understanding NASH** _____ **5**
- **NASH is progressive** _____ **5**

ABOUT REZDIFFRA (RESMETIROM)

- **About Rezdiffra** _____ **7**
- **How Rezdiffra can help you** _____ **7**
- **How Rezdiffra works** _____ **9**
- **Possible side effects** _____ **9**
- **Taking Rezdiffra** _____ **11**

SAVINGS & SUPPORT

- **Getting Rezdiffra** _____ **13**
- **Madrigal support options** _____ **13**
- **Additional NASH resources** _____ **15**
- **Have questions?** _____ **17**

IMPORTANT SAFETY INFORMATION _____ **19**

IMPORTANT SAFETY INFORMATION (cont.)

Before you take Rezdiffra, tell your healthcare provider about all of your medical conditions, including if you:

- have gallbladder problems or have been told you have gallbladder problems, including gallstones.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.

Understanding NASH

Did you know that about 1 in 5 people with fatty liver disease have **NASH**? That's why it's important to understand what NASH is and what you can do to manage it.

NASH, or nonalcoholic steatohepatitis, is a serious form of fatty liver disease. It occurs when the fat in the liver becomes toxic to you, damaging liver cells and making the liver inflamed. As the liver reacts to these changes, it gets scarred. This scarring is called fibrosis. Here are 2 things you can do to take on NASH:


-  Have your liver scarring regularly checked by your liver specialist
-  Talk to your liver specialist about treating NASH



90% of adults with NASH don't know they have it. That's why it's important to talk to your liver specialist about NASH before it causes additional liver damage.

NASH is progressive

Most people with NASH don't have any symptoms in the early stages. If it goes undetected and unmanaged for too long, **NASH can get worse and cause more serious health problems such as:**



CIRRHOSIS

~1 in 5 people with NASH and advanced scarring will develop cirrhosis in as little as ~2 years.



NEED FOR TRANSPLANT

NASH is the leading cause of liver transplants in women, and the second leading cause in men.



LIVER CANCER

~1 in 8 people with NASH and cirrhosis will develop liver cancer in less than 4 years—but liver cancer can also occur in people who have NASH without cirrhosis.



SHORTENED LIFE SPAN


NASH may shorten life expectancy.




NASH is progressive, which can lead to more serious health issues.

About Rezdiffra

A breakthrough treatment for adults with NASH with liver scarring,* **Rezdiffra is a prescription pill taken once per day and is a medicine proven to help:**



**REDUCE SCARRING*
(FIBROSIS)**



REDUCE NASH*

*Moderate to advanced liver scarring (fibrosis) without cirrhosis.

[†]Reduction in scarring defined as improvement in fibrosis with no worsening of steatohepatitis (defined as no increase in score for ballooning, inflammation, or steatosis).

[‡]Defined as resolution of steatohepatitis (score of 0-1 for inflammation, 0 for ballooning, and any value for steatosis) with no worsening of liver scarring.

IMPORTANT SAFETY INFORMATION (cont.)

Before you take Rezdiffra, tell your healthcare provider about all of your medical conditions, including if you: (cont.)

- are pregnant or plan to become pregnant. It is not

known if Rezdiffra will harm your unborn baby.

- are breastfeeding or plan to breastfeed. It is not known if Rezdiffra passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take Rezdiffra.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.

How Rezdiffra can help you

Rezdiffra was studied in a large clinical trial and showed significant results at 1 year.[‡]

Helps reduce liver scarring[§]

1.5-~2x

as many patients taking either 80 mg or 100 mg of Rezdiffra compared to placebo

Helps reduce NASH^{||}

2-~3x

as many patients taking either 80 mg or 100 mg of Rezdiffra compared to placebo

On average, liver enzymes appeared to be reduced in patients taking Rezdiffra compared to those taking placebo starting **as early as 3 months** through 1 year.

[‡]Rezdiffra was assessed by 2 credible healthcare providers (A&B):

[§]Improvement in liver scarring in patients taking Rezdiffra vs placebo:

A: 28% (100 mg of Rezdiffra), 23% (80 mg of Rezdiffra) vs 15% (placebo)

B: 24% (100 mg of Rezdiffra), 23% (80 mg of Rezdiffra) vs 13% (placebo)

^{||}NASH resolution in patients taking Rezdiffra vs placebo:

A: 36% (100 mg of Rezdiffra), 27% (80 mg of Rezdiffra) vs 13% (placebo)

B: 24% (100 mg of Rezdiffra), 26% (80 mg of Rezdiffra) vs 9% (placebo)

IMPORTANT SAFETY INFORMATION (cont.)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

- Rezdiffra and other medicines may affect each other, causing side effects. Rezdiffra may affect the way other medicines work, and other medicines may affect how Rezdiffra works.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.

How Rezdiffra works

Rezdiffra treats NASH with liver scarring* by:



WORKING DIRECTLY IN THE LIVER

Rezdiffra is specially designed to target your liver.



HELPING REDUCE LIVER FAT

Rezdiffra reactivates liver cells, allowing your liver to reduce liver fat.

*Moderate to advanced liver scarring (fibrosis) without cirrhosis.



How do I know if Rezdiffra is working?

Your liver specialist is the best person to ask. Regular visits per your specialist's recommendations can help assess the effect of Rezdiffra on your liver.

Possible side effects

Rezdiffra may cause serious side effects. Stop taking Rezdiffra and call your doctor if you develop symptoms of:

- **Liver injury (1 case in clinical trials):** including tiredness, nausea, vomiting, fever, rash, jaundice, or stomach pain/tenderness
- **Gallstones or gallbladder/pancreas inflammation:** including nausea, vomiting, fever, or stomach pain

Common side effects observed in studies include diarrhea, nausea, itching, stomach pain, vomiting, dizziness, and constipation.

See Important Safety Information on pages 18-19 for complete list of side effects.

Diarrhea and nausea were the most common side effects. Both typically began early in treatment and were mild to moderate, lasting ~3 to 4 weeks.[†]

[†]Should these problems not resolve or should they become worse, please contact your doctor.

IMPORTANT SAFETY INFORMATION (cont.)

- Especially tell your healthcare provider if you take medicines that contain gemfibrozil to help lower

your triglycerides, or cyclosporine to suppress your immune system, because Rezdiffra is not recommended in patients taking these medicines.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.



Taking Rezdiffra

Rezdiffra is a once-daily pill that you should take as prescribed for as long as your liver specialist tells you to. Here are 3 tips for taking Rezdiffra:



Take Rezdiffra
once a day, every day,
with or without food.



Keep taking Rezdiffra
as prescribed for as long as
your liver specialist tells you to.



Fit Rezdiffra into
your routine by
setting reminders.



Quick tip:

Take Rezdiffra at the same time every day, like before or after you walk the dog.

IMPORTANT SAFETY INFORMATION (cont.)

- Tell your healthcare provider if you are taking medicines such as clopidogrel to thin your blood or statin medicines to help lower your cholesterol.
- Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.

Getting Rezdiffra

Rezdiffra is sent through a specialty pharmacy in Madrigal's distribution network or through the Madrigal Patient Support program. **Here's how it works:**

1

Your liver specialist will send your prescription to a specialty pharmacy

2

The specialty pharmacy will confirm insurance coverage and out-of-pocket cost

3

The specialty pharmacy will call you to schedule your Rezdiffra shipment


4

When it's time for a refill, the specialty pharmacy will give you a call



Be sure to check your phone. Someone from your specialty pharmacy or Madrigal Patient Support will call you to set up the delivery of Rezdiffra—this may come from a number you don't recognize.

Madrigal support options



Not an actual card.

SAVE ON REZDIFFRA

Eligible, commercially insured patients may pay as little as \$10 with the Rezdiffra copay card.*



MADRIGAL PATIENT SUPPORT

Our support program is designed to help you navigate the process of accessing your Rezdiffra prescription.



Scan or visit Rezdiffra.com/patient-support to sign up for your copay card and to learn more about the Madrigal Patient Support program.

*Eligibility rules apply. Offer not valid for patients enrolled in Medicare, Medicaid, or other federal or state healthcare programs. For program terms, conditions, and eligibility criteria, visit copay.rezdiffra.com.

IMPORTANT SAFETY INFORMATION (cont.)

What are the possible side effects of Rezdiffra?

Rezdiffra may cause serious side effects, including:

- **liver injury (hepatotoxicity).** Stop taking Rezdiffra and call your healthcare provider right away if you develop

the following signs or symptoms of hepatotoxicity, including tiredness, nausea, vomiting, fever, rash, your skin or the white part of your eyes turns yellow (jaundice), or stomach pain/tenderness.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.

Additional NASH resources

If you feel like NASH is overwhelming, ask your liver specialist to recommend a local support group. You can also connect with an advocacy group dedicated to helping people with liver diseases, including NASH, such as the ones below.*



Working with advocates, policymakers, clinicians, and industry to drive the liver health field forward.



NASH knowledge

Raising public awareness of fatty liver disease and NASH by providing support, education, and resources.



Helping to improve the lives of those with fatty liver disease through awareness, screening, education, and outreach.

*The above resources are not controlled, owned, or endorsed, by Madrigal, and Madrigal is not responsible for their content.



For more support and information about Rezdiffra, visit Rezdiffra.com.



IMPORTANT SAFETY INFORMATION (cont.)

Rezdiffra may cause serious side effects, including: (cont.)

- **gallbladder problems.** Gallbladder problems such as gallstones, or inflammation of the gallbladder, or inflammation of the pancreas from gallstones can occur with NASH and may occur if you take Rezdiffra.

Call your healthcare provider right away if you develop any signs or symptoms of these conditions, including nausea, vomiting, fever, or pain in your stomach area (abdomen) that is severe and will not go away. The pain may be felt going from your abdomen to your back and the pain may happen with or without vomiting.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.

Have questions? We have answers.

When you're managing NASH with liver scarring* and starting on a new treatment, questions will pop up. We can help you find answers to questions around the topics below:

- NASH and disease progression
- About Rezdiffra
- Starting on Rezdiffra
- Side effects
- Support and assistance for Rezdiffra



Learn more about these topics.
Find answers to your questions
about Rezdiffra and NASH at
Rezdiffra.com/FAQs.

*Moderate to advanced liver scarring (fibrosis) without cirrhosis.

Be your own advocate.

Don't be afraid to ask your liver specialist questions about Rezdiffra and NASH with liver scarring.

To get the conversation started, download this discussion guide at Rezdiffra.com/guide.



Questions? Write them down!

IMPORTANT SAFETY INFORMATION (cont.)

The most common side effects of Rezdiffra include:

diarrhea, nausea, itching, stomach pain, vomiting, dizziness, and constipation.

These are not all the possible side effects of Rezdiffra. For more information, ask your healthcare provider

or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at **1-800-FDA-1088** or www.fda.gov/medwatch. You may also report side effects to Madrigal Pharmaceuticals, Inc. at 1-800-905-0324.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.

INDICATION AND IMPORTANT SAFETY INFORMATION

What is Rezdiffra?

Rezdiffra is a prescribed medicine used along with diet and exercise to treat adults with nonalcoholic steatohepatitis (NASH) with moderate to advanced liver scarring (fibrosis), but not with cirrhosis of the liver.

This indication is approved based on improvement of NASH and liver scarring (fibrosis). There are ongoing studies to confirm the clinical benefit of Rezdiffra.

Before you take Rezdiffra, tell your healthcare provider about all of your medical conditions, including if you:

- have any liver problems other than NASH.
- have gallbladder problems or have been told you have gallbladder problems, including gallstones.
- are pregnant or plan to become pregnant. It is not known if Rezdiffra will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if Rezdiffra passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you take Rezdiffra.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

- Rezdiffra and other medicines may affect each other,

causing side effects. Rezdiffra may affect the way other medicines work, and other medicines may affect how Rezdiffra works.

- Especially tell your healthcare provider if you take medicines that contain gemfibrozil to help lower your triglycerides, or cyclosporine to suppress your immune system, because Rezdiffra is not recommended in patients taking these medicines.
- Tell your healthcare provider if you are taking medicines such as clopidogrel to thin your blood or statin medicines to help lower your cholesterol.
- Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

What are the possible side effects of Rezdiffra?

Rezdiffra may cause serious side effects, including:

- **liver injury (hepatotoxicity).** Stop taking Rezdiffra and call your healthcare provider right away if you develop the following signs or symptoms of hepatotoxicity, including tiredness, nausea, vomiting, fever, rash, your skin or the white part of your eyes turns yellow (jaundice), or stomach pain/tenderness.

IMPORTANT SAFETY INFORMATION (cont.)

- **gallbladder problems.** Gallbladder problems such as gallstones, or inflammation of the gallbladder, or inflammation of the pancreas from gallstones can occur with NASH and may occur if you take Rezdiffra. Call your healthcare provider right away if you develop any signs or symptoms of these conditions, including nausea, vomiting, fever, or pain in your stomach area (abdomen) that is severe and will not go away. The pain may be felt going from your abdomen to your back and the pain may happen with or without vomiting.

The most common side effects of Rezdiffra include: diarrhea, nausea, itching, stomach pain, vomiting, dizziness, and constipation.

These are not all the possible side effects of Rezdiffra. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at **1-800-FDA-1088** or www.fda.gov/medwatch. You may also report side effects to Madrigal Pharmaceuticals, Inc. at 1-800-905-0324.

Please see the accompanying full [Prescribing Information](#), including [Patient Information](#), for Rezdiffra.



Ask about the **FIRST**
FDA-approved treatment for adults
with NASH with liver scarring,*
along with diet and exercise.
Rezdiffra.com

*Moderate to advanced liver scarring
(fibrosis) without cirrhosis.



Rezdiffra™
resmetirom tablets
60mg · 80mg · 100mg

We're here for you.

Get the latest information about Rezdiffra plus
ongoing support and resources, right in your inbox.

Scan to sign up or visit Rezdiffra.com/signup.

